



Optimization of Cultural Conditions for Enhanced Lipid Accumulation by a Local Isolate of *Cunninghamella* sp. using Response Surface Methodology

Khadiga Ali AL-Foothih

Department of Applied Microbiology, Faculty of
Applied Science, Taiz University, Yemen

Dr. Najeeb Kaid Al-Shorjani

Department of Applied Microbiology, Faculty of
Applied Science, Taiz University, Yemen

nagnas2010@gmail.com

Prof. Dr. Fahd Abdelhameed Alsharjabi

Department of Applied Microbiology, Faculty of

Received: 11 /11/2024

Accepted: 2/10/2024

Journal Website:

<https://journal.alsaeeduni.edu.ye>

تحسين الظروف المزرعية لتحسين انتاج الدهون بواسطة عزلة محلية من فطر الكانينجهاميليا باستخدام منهجية الاستجابة السطحية

الباحثة / خديجة علي محمد احمد الفتيح

قسم الميكروبيولوجي، كلية العلوم التطبيقية

جامعة تعز، اليمن

د/ نجيب قائد الشرجاني

قسم الميكروبيولوجي، كلية العلوم التطبيقية

جامعة تعز، اليمن

أ.د/ فهد عبد الحميد الشرجبي

قسم الميكروبيولوجي، كلية العلوم التطبيقية

جامعة تعز، اليمن

الملخص

يعد إنتاج الدهون الميكروبية بواسطة الفطريات الدهنية مصدرًا واعدًا لإنتاج الأحماض الدهنية المتعددة غير المشبعة، والتي تعتبر مواد ذات قيمة للاستخدامات الغذائية والصيدلانية، ويعد تحسين ظروف البيئية والزراعية أمرًا ضروريًا، لتعزيز إنتاج الدهون الميكروبية. يهدف هذا البحث إلى تحسين إنتاج الدهون باستخدام عزلة محلية من فطر الكانينجهاميليا، حيث تم دراسة تأثير عدة عوامل مثل مصدر الكربون، ومصدر النيتروجين، ودرجة الحموضة، ووقت التحضين على تراكم وإنتاج الدهون بواسطة فطر الكانينجهاميليا بطريقة تقليدية تعتمد على تغيير عامل واحد في كل مرة، أظهرت النتائج أن أفضل مصدر للكربون كان الجلوكوز، وأفضل مصدر للنيتروجين كان نترات الصوديوم. كما وجد أن القيمة المثلى للأس الهيدروجيني (درجة الحموضة pH) هي 6.0 وأن أفضل وقت للتحضين هو 5 أيام، بالإضافة إلى ذلك، تم تحسين تركيز الجلوكوز، ونواتج الصوديوم، ودرجة الحموضة، لتحسين إنتاج الدهون باستخدام منهجية الاستجابة السطحية (RSM)، وتم تطبيق تصميم مركب مركزي (CCD)، واستخدام نموذج الانحدار متعدد الحدود مع الحد التربيعي لتحليل البيانات التجريبية باستخدام تحليل التباين (ANOVA)، وأظهرت نتائج التحليل أن التركيز الأمثل للجلوكوز كان 38.28 جم/لتر، ونواتج الصوديوم 0.48 جم/لتر، وذلك عند قيمة pH بلغت 5.79، مما أدى إلى تراكم للدهون بنسبة 25.4% (وزن/وزن)، وكشف النموذج التربيعي أن الأس الهيدروجيني (درجة الحموضة) كان العامل الأكثر تأثيرًا في تحسين إنتاج الدهون بواسطة فطر الكانينجهاميليا، مما يدل على قدرة هذه العزلة المحلية في تحقيق تراكم فعال لهذه الدهون. الكلمات المفتاحية: فطر كانينجهاميليا، الأحماض الدهنية العديدة غير المشبعة، الدهون الميكروبية، التحسين، منهجية الاستجابة السطحية.

Optimization of Cultural Conditions for Enhanced Lipid Accumulation by a Local Isolate of *Cunninghamella* sp. using Response Surface Methodology

Khadiga Ali AL-Footih

Department of Applied Microbiology, Faculty of Applied Science,
Taiz University, Yemen

Dr. Najeeb Kaid Al-Shorjani

Department of Applied Microbiology, Faculty of Applied Science,
Taiz University, Yemen

Prof. Dr. Fahd Abdelhameed Alsharjabi

Department of Applied Microbiology, Faculty of Applied Science,
Taiz University, Yemen

Abstract

Microbial lipid production by oleaginous fungi offers a potential promising source for producing polyunsaturated fatty acids (PUFAs), valuable compounds for nutraceutical and pharmaceutical applications. Optimization of culture conditions is essential for enhancement of microbial lipid production. The aim of this study is to improve the lipid synthesis using a local oleaginous mold of *Cunninghamella* spp. The effects of several factors including carbon source, nitrogen source, pH and incubation time on lipid accumulation by *Cunninghamella* spp were investigated conventionally (one-variable-at-a-time). The results showed that the most effective carbon source was glucose and sodium nitrate was the best nitrogen source for lipid synthesis. The optimum pH and incubation time were found to be 6.0 and 5 days, respectively. Furthermore, glucose concentration, sodium nitrate and pH were further optimized to maximize lipid production using response surface methodology (RSM). A central composite design (CCD) was applied, and a polynomial regression model with a quadratic term was used to estimate the experimental data using analysis of variance (ANOVA). Results of RSM-CCD optimization indicated that the optimal concentrations of glucose and sodium nitrate were 38.28 g/L glucose, 0.48 g/L, respectively, at a pH value of 5.79, resulting in a lipid accumulation of 25.4% (w/w). The quadratic model revealed that pH was the most influential factor in lipid synthesis by *Cunninghamella* spp., a local isolate demonstrating efficient lipid accumulation potential.

Keywords: *Cunninghamella* spp; polyunsaturated fatty acids; microbial lipids; optimization; response surface methodology.

Introduction:

Lipids, a major macronutrient category encompassing oils and fats, play diverse roles in metabolism. They serve as an energy reserve in both animal and plant bodies, form the primary component of cell membranes, and act as precursors for biosynthesis of various hormones (Akpınar-Bayazit, 2014).

Lipids or oils derived from microorganisms are known alternatively as single-cell oils (SCOs) (Mhlongo, et al., 2021). SCOs are hydrophobic compounds primarily composed of triacylglycerols which are derived from oleaginous microorganisms such as molds, yeasts, algae and bacteria (Rossi et al., 2011). Oleaginous microorganisms are lipid-producing organisms that can accumulate oils to over 20% of the total dry cell mass within their cells or mycelia (Chebbi et al., 2019; Ratledge, and Wynn, 2002). These microorganisms are emerging as promising cell factories for the production of SCOs (Leong et al., 2018). SCOs possess significant industrial and economic value as potential to replace costly plant- or animal-based lipids. They have many applications and can be used as alternative to costly lipid sources such as fish oil, borage oil, cocoa butter, and evening primrose oil, as feedstock for second-generation biofuel production and as precursors for the synthesis of several oleochemical compounds (Kalampounias, et al., 2024; Robles-Iglesias, et al., 2023; Lewis et al., 2000). The chemical composition of lipids produced by oleaginous fungi primarily consists of unsaturated or polyunsaturated fatty acids (PUFAs). Among these PUFAs, the fatty acid gamma-linolenic acid (GLA— $\Delta^{6,9,12}$ C18:3) is found in varying concentrations within the cellular lipids of these microorganisms (Bellou et al., 2016; Papanikolaou and Aggelis, 2019). It is noteworthy that the gamma (γ) isomer of linolenic acid (GLA), rather than the more common alpha (α) isomer, is a unique characteristic of Zygomycetes (Ratledge, and Wynn, 2002; González-Fernández et al., 2020).

Lipids containing GLA are highly valuable commodities for both the nutraceutical and pharmaceutical industries due to their properties such as anti-thrombotic, anti-irritant, and particularly anti-cancer. They have also demonstrated effective in treating various diseases, including inflammatory disorders, atopic eczema, and rheumatoid arthritis (Ratledge and Wynn, 2002; Ratledge, 1994). Despite the fact that vegetable oils containing GLA are generally expensive, the potential for developing a fermentation bioprocess

to produce significant amounts of fungal oils rich in GLA from low-cost substrates offers significant promise (Bellou et al., 2016; Fazili et al., 2022; Athenaki et al., 2018).

The global demand for lipids is expanding at a faster rate than traditional production methods can meet due to the limited arable land and increase in population growth. Production of lipids by oleaginous microorganisms, with their distinctive properties and applications in the food, pharmaceutical, chemical, and energy industries, provides a promising and sustainable alternative oil source (Huang et al., 2017).

Microbial SCO production offers several benefits over traditional animal and vegetable oils. Microbial sources have shorter life cycles, exhibit rapid growth rates, are easier to cultivate in bioreactors, are readily scalable, and are unaffected by climate changes (Li et al., 2008) which make them ideal candidates for industrial-scale production (Dong, 2016).

Cunninghamella species are oleaginous fungi, belong to Zygomycetes, capable of synthesizing and accumulating polyunsaturated fatty acids (PUFAs). Among PUFAs, the bioactive GLA, which has potential anti-tumor activities, and docosahexaenoic acid (DHA), which is essential for human health, especially for brain and eye development, are particularly noteworthy (Kalampounias et al., 2024; Manikan et al., 2015).

In order to improve the productivity and yield of microbial lipids through fermentation processes, culture conditions and nutritional factors must be optimized. Lipid accumulation by oleaginous microorganisms and fatty acid content differ depending on the environmental conditions such as pH, incubation time, incubation temperature and the kind of employed microorganism as well as the type and availability of key nutrients (Ageitos et al., 2011)

Fermentation optimization can be achieved using a "one variable at a time" method to determine the optimal conditions for maximizing growth and lipid synthesis. However, optimization using response surface methodology (RSM) has been used much more frequently in recent years to optimize medium content and culture conditions to improve the efficacy of fermentation process (Awad et al., 2011). When compared to traditional optimization methods, RSM is a highly helpful technique that minimizes the number of tests, saving time and chemicals utilized in the experiments. RSM

also yields accurate response predictions, which makes it a favorable choice for experimental design (Singh et al., 2013). The main designs in RSM include the Box–Behnken design (BBD), central composite design (CCD), user-defined design, D-Optimal design, historical data design and one-factor design. The BBD and CCD are the two most popular RSM statistical techniques and BBD only offers three levels for a single numerical variable, compared to five for CCD (Al-Shorgani et al., 2015 and Bezerra et al., 2008). Additionally, RSM offers an experimental model that forecasts the connections and interplay between a collection of experimental variables and observed results enabling the optimization of experimental setting (Zheng et al., 2008).

The objective of the present study was to optimize lipid synthesis by RSM with CCD and analyzing the impact of glucose concentration, sodium nitrate concentration and pH as well as their interactions on lipid synthesis using a local fungal isolate of *Cunninghamella* spp

MATERIALS & METHODS:

Microorganism Strain:

Cunninghamella spp is a newly local fungal isolate which was recovered from a local soil from Taiz City. *Cunninghamella* spp was presumptively identified according to its morphology by light microscope. This fungal isolate was stored in potato dextrose agar (PDA) slant agar at 4°C.

Inoculum Preparation and Culture Conditions:

Fresh inoculum of *Cunninghamella* spp was prepared by transferring mycelium from PDA into a fresh potato dextrose broth (PDB) and incubated at 28°C for 48 hours. This fresh inoculum was then used to inoculate the production medium.

Lipid production was conducted in batch fermentation using flask scale (250 mL conical flasks) with a working volume of 100 mL. The production medium used for lipid production was basal medium. *Cunninghamella* spp was grown on basal medium that contains 50 g/L glucose, 0.5 g/L yeast extract, 0.4 g/L $MgSO_4 \cdot 7H_2O$, 2 g/L KH_2PO_4 , 0.5 g/L $CaCl_2$, 0.05 g/L $CuSO_4 \cdot 5H_2O$, with an initial pH of 6.0. A 10% (v/v) of fresh mycelium from PDB was added as inoculum to 100 mL of basal media in 250 mL conical flasks and the mixture was then incubated under constant conditions for five days at 28°C. After the incubation period, the fungal culture was extracted

using filtration and the biomass was collected and repeatedly washed using purified water to eliminate any remaining medium (Youssef et al., 2020).

Verification of the Quadratic Model:

The predicted optimum conditions by the empirical model were validated using a set of batch culture fermentation under optimum conditions of glucose, sodium nitrate and initial pH.

Dry Cell Weight Determination:

Fungal biomass were harvested by centrifugation at 3000 rpm for 20 minutes followed by two rinsing the biomass pellet twice with 50 mL sterile distilled water. Samples were dried in hot air oven at 95 °C until reach constant weights. Biomass was expressed as dried weight in gram per liter of growth medium.

Lipid Extraction:

Extraction of lipid was conducted according to Folch et al., (1957). After drying the mycelium biomass of *Cunninghamella* spp., the dried biomass was then crushed into a powder using a mortar and pestle, and was added to a mixture of 20 mL chloroform and 10 mL methanol and the mixture was stirred for 30 minutes. Filtration by filter paper (Wattmann No.1) was used to separate the mixtures and sodium chloride solution 0.9% was added. Lipids-containing solvent was isolated using a separating funnel. The layer on the bottom of NaCl, methanol, and water were eliminated, and the residual of solvent was dried and then weighted. The lipid content was estimated by using Equation (1) according to Nisha et al., (2009) as follow:

$$\text{Lipid content (\%)} = \frac{\text{Weight of lipid (g)}}{\text{Weight of dried biomass (g)}} \times 100 \quad (\text{Equation 1})$$

Experimental Design of Culture Conditions for Lipid Synthesis Optimization by *Cunninghamella* spp:

Optimization of culture conditions influencing the lipid biosynthesis by *Cunninghamella* spp were estimated in two steps, preliminary optimization using covenantal method of one-variable-at-a-time (OVAT) and final optimization (secondary) using RSM with CCD.

Preliminary optimization for lipid production by a local isolate of *Cunninghamella* spp was conducted to evaluate the impact of various parameters including various carbon sources, nitrogen sources and initial pH and incubation time using (OVAT) method. The tested carbon sources include glucose, fructose, lactose, dextrose, maltose, sucrose and starch. To

choose the best nitrogen source that give the highest lipid production, yeast extract (the original nitrogen source in basal medium) was replaced with three other nitrogen sources, one at a time, on equal nitrogen bases (0.5 g/L); sodium nitrate, peptone, and ammonium acetate. The influence of initial pH value of the fermentation medium on fungal biomass growth and lipid accumulation was studied which ranged from pH 4 to 9. Incubation time of the fermentation culture was varied from 4 days to 7 days in order to estimate the suitable period for maximal lipid production by local strain of *Cunninghamella* spp.

Secondary optimization for lipid production was carried out based on the results obtained from the preliminary optimization. Three variable parameters including glucose concentration, sodium nitrate concentration and initial pH were chosen at two levels utilizing the design of experiment (DOE) by RSM based on CCD. The chosen ranges of the three variables are as follow: glucose concentration 20 – 60 g/L, sodium nitrate concentration 0 – 1 g/L and initial pH of the medium 4 – 9. Table (1) shows the ranges of the tested three variables.

Table (1): The level of variable parameters affecting lipid production by *Cunninghamella* spp

Parameter	Unite	Level (-1)	Level (+1)
Glucose (X_1)	g/L	20	60
Sodium nitrate (X_2)	g/L	0	1
pH (X_3)	-	4	9

Experimental design by RSM-CCD with six-center point replications gave 20 experimental runs which were conducted with various combinations of three independent variables (Table (2)). These experiments took place in batch fermentation cultures of *Cunninghamella* spp within 250 mL conical flasks, each containing a working volume of 100 mL. The inoculum size was 10% for each culture and the incubation temperature was set at 28°C for 5 days incubation time. Statistical analysis, performed by an Analysis of Variance (ANOVA) with DOE software was employed to determine the best culture conditions for the maximum lipid production by the strain of *Cunninghamella* spp.

Table (2): Experimental design matrix using RSM-CCD for three variables for lipid optimization by *Cunninghamella* spp

Run No.	X ₁ : Glucose (g/L)	X ₂ : Sodium nitrate (g/L)	X ₂ : pH
1	60.00	0.05	9.0
2	60.00	1.00	4.0
3	40.00	0.53	2.2
4	40.00	0.53	6.5
5	20.00	1.00	9.0
6	40.00	0.53	6.5
7	40.00	0.53	6.5
8	60.00	0.05	4.0
9	20.00	1.00	4.0
10	40.00	0.53	6.5
11	73.64	0.53	6.5
12	40.00	0.53	6.5
13	40.00	0.00	6.5
14	6.36	0.53	6.5
15	20.00	0.05	4.0
16	40.00	0.53	10.7
17	60.00	1.00	9.0
18	40.00	0.53	6.5
19	20.00	0.50	9.0
20	40.00	1.32	6.5

Results and Discussion

Previous studies have identified that incubation temperature, incubation period, initial pH value and the sources of carbon and nitrogen play as crucial factors effecting lipid synthesis by oleaginous microorganisms (Enshaeieh et al., 2014). In this investigation, the initial optimization step utilized a "one variable at a time" experimental method to determine the major parameters necessary for maximizing lipid synthesis by a local isolate of *Cunninghamella* spp. The preliminary optimization of lipid production utilizing OVAT used

to optimize carbon source, nitrogen source, initial medium pH and incubation time.

Impact of Different Carbon Sources on Lipid Synthesis by *Cunninghamella* spp:

The type of carbon source in a fermentation medium significantly affects the amount and the composition of lipids produced by various oleaginous fungal species, making it as an essential medium component for fungal growth. This is due to their unique metabolic processes.

Glucose is the most common sugar often utilized by oleaginous fungi for biomass growth and lipid accumulation under nitrogen-limited conditions (Carvalho et al., 2018). Furthermore, the concentration of glucose in the culture medium was another key factor that boosted lipid production by oleaginous molds. The optimal glucose concentration for achieving maximal lipid accumulation in oleaginous molds is varied within the range of 20 to 80 g/L (Lei et al., 2024). Al-Hawash et al., (2018) found that presence of low glucose concentrations in culture medium resulted in lower lipid yield, whereas rising the amounts of glucose boosts lipid content, peaking at approximately 50g/L with a lipid synthesis content of 39.02% of biomass.

The impact of various carbon sources on lipid production by *Cunninghamella* spp was investigated in this study. The tested carbon sources included monosaccharides, disaccharides, and polysaccharides such as glucose, fructose, lactose, maltose, sucrose, dextrose and starch. The experiments were conducted using a basal medium, inoculum amount of 10% (v/v), incubation temperature of 28°C, incubation time of 7 days and under static condition.

Results in this study demonstrated that the lipid biosynthesis and fungal biomass growth by *Cunninghamella* spp was significantly influenced by the carbon type supplied in the culture medium. It was observed that all tested carbon sources were consumed by the isolate *Cunninghamella* spp while the biomass growth and lipid content were varied based on the carbon source used. Among the tested carbon sources, glucose yielded the highest lipid accumulation, followed by sucrose, as illustrated in Figure (1). Starch, on the other hand, resulted in the lowest fungal growth and lipid production. This can be attributed to the fact that glucose is a simple carbon source which is more easily metabolized by the fungus *Cunninghamella* spp. As a result, it

leads to greater lipids and biomass production compared to more complex carbon sources such as starch (Nasr et al., 2017; Nisha and Venkateswaran, 2011).




Figure (1): Impact of various carbon sources on lipid synthesis by *Cunninghamella* spp

Several studies have reported that glucose yielded the highest lipid content among various carbon sources using oleaginous mold *Cunninghamella* spp (Ahmed et al., 2006; Papanikolaou et al., 2007; Shrivastava et al., 2008; Sukrutha et al., 2014). A study by Varma et al., (2018) to investigate the effect of carbon sources on the synthesis of γ -linolenic acid (GLA) using *Cunninghamella elegans* CFR C07 demonstrated that glucose was the most effective carbon source for synthesis of GLA lipid. Our findings are consistent with these studies, indicating that the best source of carbon for synthesis of lipid using a local isolate of *Cunninghamella* spp is glucose. Glucose as a common carbon source is known to improve both growth and lipid synthesis in oleaginous molds (Ochsenreither et al., 2016).

Impact of Different Nitrogen Sources On Lipid Synthesis by *Cunninghamella* spp:

Depletion of nitrogen source is concerned as the most influencing parameter for lipogenesis in majority oleaginous fungal species. Initially, when there is an adequate amount of nitrogen source, oleaginous fungi utilize the available source of carbon in the medium for growth and cell proliferation, producing lipid-free biomass that contains functional lipids. Once the

nitrogen in the culture medium is exhausted in the presence of accessible carbon, oleaginous fungi then initiate the accumulation and storage of lipids (Economou et al., 2011).

In this study, the impact of different nitrogen sources on lipid biosynthesis by *Cunninghamella* spp was investigated. This was conducted using four different nitrogen sources in a basal medium contains glucose (50 g/l) as the sole carbon source under optimal conditions of incubation temperature at 28°C, initial pH of 6.0 and under static condition. The results of this study indicated that sodium nitrate was the optimal inorganic nitrogen source for maximum lipid content (22.7%) followed by yeast extract (Figure (2)). In terms of biomass growth and lipid production, yeast extract showed the highest fungal growth as 3.3 g/L, and the produced lipid weight was 0.7 g/L, whereas the lipid content was 21.2% as illustrated in Figure (2). In contrast, the lowest lipid yield, in terms of lipid/dry biomass ratio, was observed with peptone and ammonium acetate.

The results in this study are in agreement with that previously reported by Sheekh et al., (2019) who reported that addition of sodium nitrate to the growth medium of microalga *Desmodesmus intermedius* enhanced lipid production by 60% compared to the control. Similarly, Abdella et al., (2020) found that sodium nitrate produced the highest lipid production by *Aspergillus* spp with 53.3% lipid/dry biomass, followed by ammonium sulfate (45.3%) and ammonium oxalate (39.6%) while the lowest lipid yield was obtained with ammonium chloride as 34.3% lipid/dry biomass. In addition, Moubasher et al., (2016) demonstrated that supplementing the growth medium of *Fusarium oxysporum* with sodium nitrate as a nitrogen source led to maximize lipid accumulation (23.65%) and biomass (2.10 g/L). Ali et al., (2017) reported that utilization of sodium nitrate as a nitrogen source for lipid production by *Penicillium brevicompactum* resulted in the highest lipid accumulation. In contrast, Ramírez-Castrillón et al., (2017) and Xing et al., (2012), reported different findings, showing that the most appropriate nitrogen source for lipid production was ammonium sulfate using *Mortierella isabellina* and *Meyerozyma guilliermondii*, respectively.




Figure (2): Impact of nitrogen source on lipid synthesis by *Cunninghamella* spp.

Impact of Initial pH on Lipid Synthesis by *Cunninghamella* spp:

The influence of initial medium pH on lipid synthesis by oleaginous *Cunninghamella* spp was studied over a range between pH 4.0 to pH 9.0 by adjusting the medium's pH before autoclaving by dilute HCl/NaOH. The results in this study found that the lipid production was maximized when *Cunninghamella* spp was cultured at pH 6.0, resulting in the highest biomass growth and lipid content. The results also indicated that the total lipid content extremely minimize at pH higher than 8.0 (Figure(3)). This observation is supported by Sukratha et al., (2014) who reported that the highest lipid accumulation by *Cunninghamella blakesleeana* was reached at pH 6.0. Moreover, Shoaib et al., (2014) indicated that pH 6.0 was the optimal for the growth and lipid production by *Aspergillus wentii*. Additionally, this outcome is close to the optimal pH of 5.5 for lipid production by *Aspergillus awamori* as reported by Venkata and Venkata (2014).




Figure (3): Impact pH on lipid synthesis by *Cunninghamella* spp.

Studies have shown that a pH range of 5.0 to 6.0 is the optimal pH for the highest lipid accumulation and growth of the majority of oleaginous molds (Ali and El-Ghonemy 2014; Ruan et al., 2014; Jiru et al., 2017; Ali et al., 2017). Previous studies have indicated that pH is a critical environmental factor that can induce stress responses in fungal cells, subsequently influencing their growth, lipid accumulation, and the composition of the lipids produced (Xia et al., 2011; Mironov et al., 2018).

Impact of Incubation Period on Lipid Synthesis by *Cunninghamella* spp:

The incubation time significantly influences lipid accumulation in oleaginous fungal species and the optimal incubation durations varying depends on the fungal species, fermentation medium and growth conditions (Nisha and Venkateswaran, 2011).

The current study found that on the fifth day of growth, there was the greatest biomass growth and lipid synthesis (Figure (4)). The incubation time influenced fungal biomass formation and lipid synthesis, as a longer incubation period allowed for more effective use of the obtainable source of carbon in the growth culture, leading to greater lipid content. Shorter incubation periods resulted in less substrate utilization by *Cunninghamella* spp, resulting in reduced biomass formation and lipid accumulation. These results are consistent with findings by Kumar and Banerjee (2013) who

reported that most lipid synthesis by *Aspergillus* spp occurred later five days of incubation.

In addition, Ali and El-Ghonemy (2014) noted that both *Trichoderma viride* and *Aspergillus* spp reached most lipid synthesis after five days of cultivation. Abdelhamid et al., (2019) also indicated that *Penicillium commune* reached the highest lipid synthesis using a basal liquid medium after 5 days of incubation.

The maximum lipid synthesis by *Cunninghamella* spp was found to occur after 5 days of cultivation under constant conditions at 28°C and pH of 6.0, based on the findings from the first step of sequential optimization. Moreover, the best sources of carbon and nitrogen were found to be glucose and sodium nitrate, respectively.




Figure (4): Impact of incubation period on lipid synthesis by *Cunninghamella* spp.

Optimization of Glucose, Sodium nitrate and pH using RSM:

In order to optimize the quantities of glucose and sodium nitrate in the fermentation medium, and the initial medium' pH were further optimized using RSM-CCD, to enhance lipid production by this local isolate of *Cunninghamella* spp.

RSM has been effectively employed for optimization of several bioproducts fermentation (Lin et al., 2011; Ranjan et al., 2013b). In this

regards, the impact and the interaction between three experimental variable factors (glucose, sodium nitrate, and pH) were determined to enhance and maximize lipid production by *Cunninghamella* spp. using RSM-CCD. Experimental design by RSM-CCD was employed, resulting in 20 experimental tests. After conducting these experiments, the lipid production content was ranged from 0.6% to 25%, as shown in Table (3).

Table (3): Experimental design for three experimental factors and results of lipid accumulation by *Cunninghamella* spp.

Run No.	Variable parameters			Response
	X ₁ : Glucose (g/L)	X ₂ : Sodium nitrate (g/L)	X ₃ : pH	Lipid content (%)
1	60.00	0.05	9.0	0.9
2	60.00	1.00	4.0	10
3	40.00	0.53	2.2	0.6
4	40.00	0.53	6.5	22
5	20.00	1.00	9.0	2.5
6	40.00	0.53	6.5	21
7	40.00	0.53	6.5	24.2
8	60.00	0.05	4.0	11.5
9	20.00	1.00	4.0	3.4
10	40.00	0.53	6.5	22.5
11	73.64	0.53	6.5	4.8
12	40.00	0.53	6.5	25
13	40.00	0.00	6.5	15
14	6.36	0.53	6.5	0.8
15	20.00	0.05	4.0	1.5
16	40.00	0.53	10.7	0.0
17	60.00	1.00	9.0	0.6
18	40.00	0.53	6.5	23.8
19	20.00	0.50	9.0	1.0
20	40.00	1.32	6.5	10

The results were analyzed using one-way ANOVA (analysis of variance) suitable for the experimental design, as shown in Table(4). The ANOVA analysis of the quadratic regression model indicates that the model is significant ($p < 0.0001$).

The Model F-value of 48.8 suggests a high level of significance, as it is the ratio of the mean square regression to the mean square residual. Additionally, the Model P-value (Prob > F) is very low (0.0001), further confirming the model's significance.

Table (4): Analysis of variance (ANOVA) for the quadratic polynomial model of lipid accumulation by *Cunninghamella* spp.

p-value (Prob > F)	F value	Mean ²	DF	Sum of Squares	Source
< 0.0001	48.81932	194.4401	9	1749.961	Model
0.0161	8.362222	33.30549	1	33.30549	Glucose (X ₁)
0.5459	0.390826	1.556602	1	1.556602	Sodium nitrate (X ₂)
0.0125	9.232155	36.7703	1	36.7703	pH (X ₃)
0.3786	0.848638	3.38	1	3.38	X ₁ X ₂
0.0081	10.8578	43.245	1	43.245	X ₁ X ₃
0.8901	0.020086	0.08	1	0.08	X ₂ X ₃
< 0.0001	174.0008	693.0196	1	693.0196	X ₁ ²
< 0.0001	51.47483	205.0166	1	205.0166	X ₂ ²
< 0.0001	221.4023	881.8126	1	881.8126	X ₃ ²
		3.982852	10	39.82852	Residual
0.1658	2.52829	5.708037	5	28.54018	Lack of Fit
		2.257667	5	11.28833	Pure Error
			19	1789.79	Cor Total
				Mean = 10.06	Std. Dev.= 2.0
Adequate Precision = 17.5				Adj R ² = 0.95	R ² = 0.97

To determine the significance of each variable factor and consequently, the pattern of the reciprocal interactions among the test variables, the p-values were employed as a tool. Table (4) provides the F-value, accompanying p-values, and the coefficient estimate. The model terms are considered significant when the value of P is less than 0.0001. The study's test variables, X₁, X₂, X₃, X₁ X₂, X₁ X₃, X₂ X₃, X₁², X₂², and X₃² (where X₁ is glucose, X₂ is

sodium nitrate and X_3 is pH), appear to be significant model terms based on the coefficient estimates and corresponding p-value. X_1 , X_3 , $X_1 X_3$, X_1^2 , X_2^2 , and X_3^2 ($P < 0.05$) have significant effect on lipid production. However, it was observed that X_2 and the interactions between $X_1 X_2$ and $X_2 X_3$ were not significant ($P > 0.05$).


ANOVA was used in a quadratic regression study to assess the model coefficients' significance. The strength of the interaction between each independent variable is also reflected in the impact of every variable, as shown through the p-values. The model in terms of real variables and lipid synthesis as the expected response is presented in Equation 2, as follow:

Lipid content (%w/w)


$$\begin{aligned} &= -75.58363 + 1.79714 \times \text{Glucose} + 24.65162 \\ &\times \text{Sodim nitrate} + 17.35919 \times \text{pH} - 0.068421 \times \text{Glucose} \\ &\times \text{Sodim nitrate} - 0.046500 \times \text{Glucose} \times \text{pH} + 0.084211 \\ &\times \text{Sodim nitrate} \times \text{pH} - 0.017261 \times \text{Glucose}^2 - 20.64178 \\ &\times \text{Sodim nitrate}^2 - 1.24613 \\ &\times \text{pH}^2 \end{aligned} \quad (\text{Equation 2})$$

The goodness of fit of the model was assessed using the coefficient of determination (R^2). The R^2 expresses the extent to which the independent variables and their interactions account for the variability observed in the experimental responses. According to Alharbi et al., (2022) the R^2 value denotes the optimal situation in which the model fully explains the variation in actual data. The study's R^2 value was 0.977, meaning that the model could account for 97.77% of the response variability. Furthermore, all regression models' adjusted R^2 value were high and nearly matched the R^2 value that were anticipated, demonstrating a remarkable agreement between the actual and expected findings. A ratio larger than four indicates adequate precision, which measures the signal-to-noise ratio (Fawzy et al., 2022). In this study, the predicted R^2 of 0.8597 was reasonably close to the adjusted R^2 of 0.9577, and the adequate precision ratio of 17.566 indicated a sufficient signal.


The primary and interaction impacts of the covariates on the response value were better understood with the use of three-dimensional response surface plots, which visually displayed the regression equation. The response surfaces of glucose and pH, glucose and sodium nitrate, and sodium nitrate and pH on lipid synthesis are shown in Figures (5a–c), respectively.



(a)



(b)



(c)

Figure (5): Response surface plot of lipid content using *Cunninghamella* spp observing the interaction between (a) impact of glucose and sodium nitrate contents on lipid synthesis, (b) impact of glucose and pH on lipid synthesis, (c) impact of sodium nitrate and pH on lipid synthesis.

Figure (5a) illustrates the influence of varying glucose and sodium nitrate concentrations on lipid synthesis. The highest lipid synthesis was achieved with concentration of glucose 50 g/L and a sodium nitrate concentration of 0.63 g/L. Increasing the glucose concentration resulted in a continuous rise in lipid synthesis.

Figure (5b) demonstrates the combined impacts of glucose and pH on lipid synthesis. The surface plot indicates that, with a constant sodium nitrate concentration of 0.53 g/L, the highest lipid synthesis occurred at a concentration of glucose 50 g/L and a pH of 6.5.

Figure (5c) shows the influence of sodium nitrate and pH on lipid synthesis. The 3D plot reveals no significant interaction between sodium nitrate and pH, as detailed in Table (4).

Model Validation:

Based on Table (3), the highest lipid content produced by *Cunninghamella* spp was 25% (w/w) was achieved with glucose, sodium nitrate, and pH levels set at 40.0 g/L, 0.53 g/L, and 6.5, respectively.

The quadratic model proposed optimal conditions for maximal lipid accumulation to be 38.28 g/L of glucose, 0.48 g/L of sodium nitrate, and a pH of 5.79, with a predicted lipid production of 23.55%. The model's validity was confirmed through validation experiments conducted in triplicates under the estimated optimal conditions. These experiments resulted in a lipid production of 25.4%, closely matching the predicted value and confirming the model's adequacy as show in Table (5).

Table (5): Model validation of optimized factors on lipid production under optimized conditions

Optimized Conditions (g/L)			Lipid Content (%)	
Glucose	Sodium nitrate	pH	Experimental	Predicted
38.28	0.48	5.79	25.4%	23.55%

Numerous studies have optimized lipid synthesis via RSM. Shrivastava et al. (2008) optimized fermentation conditions for the oleaginous fungus *Cunninghamella echinulata* var. *elegans* using a CCD, resulting in a lipid synthesis of 19.8 ± 0.35 mg/g of dry cell weight. Bardhan et al., (2019) obtained the maximum lipid synthesis of 1.73 g/L by *Penicillium citrinum* using CCD-based RSM.

Additionally, Shoaib et al. (2018) achieved a maximum lipid synthesis of 40% from *Aspergillus wentii* biomass when cultivated in a growth culture containing 50 g/L glucose, 1 g/L nitrate, and 1.5 g/L phosphate using RSM. Optimization of lipid production by oleaginous microorganisms using RSM technique exhibited that RSM-based optimization is an efficient technique for maximal lipid generation.

Conclusion:

Cunninghamella spp., a local fungal isolate, was used successfully for lipid production. Enhanced lipid synthesis was optimized through the utilization of Cunninghamella spp using RSM-CCD. The optimized conditions for the maximum lipid-enriched biomass were optimized. A maximum lipid content of 25.4 % was obtained using RSM-based CCD after 5 days of incubation at pH 5.79 and glucose, sodium nitrate concentrations of 38.28 g/L and 0.48g/L, respectively. Thus, it can be concluded that the lipid-enriched biomass was successfully optimized using a local isolate of Cunninghamella spp.

References:

- Abdelhamid, S. A.; Hussein, A. A.; Asker, M. S.; El Sayed, O. H. and Mohamed, S. S. (2019). Optimization of culture conditions for biodiesel production from Egyptian isolate *Penicillium commune* NRC2016. Bulletin of the National Research Centre. 43, 1-9.
- Abdellah, E. M.; Ali, T. H.; Abdou, D. A. M.; Hassanein, N. M.; Fadel, M.; El-Din, A. K. and El-Ghonemy, D. H. (2020). Enhancement of lipid productivity from a promising oleaginous fungus *Aspergillus* sp.strain EM2018 for biodiesel production: Optimization of culture conditions and identification. *Grasas y Aceites*. 71(3): e371-e371.
- Ageitos J. M; Vallejo J. A; Veiga-Crespo, P. and Villa T.G. (2011). Oily yeasts as oleaginous cell factories, *Appl Microbiol Biotechnol*. 90:1219–1227.
- Ahmed, S. U.; Singh, S. K.; Pandey, A.; Kanjilal, S. and Prasad, R. B. (2006). Effects of various process parameters on the production of g-linolenic acid in submerged fermentation. *Food Technol Biotechnol*. 44; 283-287.
- Akpinar-Bayizit, A. (2014). Fungal Lipids: The Biochemistry of Lipid Accumulation. *Int J Chem Eng Appl*. 5:409–414.
- Alharbi, N.K.; Al-Zaban, M.I.; Albarakaty, F.M.; Abdelwahab, S.F.; Hassan, S.H. and Fawzy, M.A. Kinetic, (2022). Isotherm and thermodynamic aspects of Zn²⁺ biosorption by *Spirulina platensis*: Optimization of process variables by response surface methodology. *Life*. 12: 585.
- Al-Hawash, AB., Li, S. Zhang, X., Zhang, X. and Ma, F. (2018). Productivity of linoleic acid by oleaginous fungus *Cunninghamella echinulata* using a pulsed high magnetic field. *Food Biosci* 21:1–7.

- Al-Shorgani, N. K. N.; Mohd Isa, M. H.; Yusoff, W. M. W.; Kalil, M. S. and Hamid, A. A. (2015). Response Surface Methodology for Biobutanol Optimization Using Locally Isolated *Clostridium acetobutylicum* YM1. *International Journal of Green Energy*. 12 (12): 1236-1243.
- Ali, T. H. and El-Ghonemy, D. H. (2014). Optimization of culture conditions for the highest lipid production from some oleaginous fungi for biodiesel preparation. *Asian Journal of Applied Sciences*. 2(5): 2321–0893.
- Ali, T. H.; El-Gamal, M. S.; El-Ghonemy, D. H.; Awad, G. E. and Tantawy, A. E. (2017). Improvement of lipid production from an oil-producing filamentous fungus, *Penicillium brevicompactum* NRC 829, through central composite statistical design. *Annals of Microbiology*. 67(9): 601-613.
- Athenaki, M., Gardeli, C., Diamantopoulou, P., Tchakouteu, S. S., Sarris, D., Philippoussis, A. and Papanikolaou, S. (2018). Lipids from yeasts and fungi: Physiology, production and analytical considerations. *J. Appl. Microbiol*. 124: 336–367.
- Awad, G. E. A.; Salam, W.A.; El-diwany, A.; Salama, B.A.; Abdelkader, A.F. and Esawy, M.A.D. (2011). Keratinase production by *Bacillus pumilus* GHD in solid-state fermentation using sugar cane bagasse: optimization of culture conditions using a box-Behnken experimental design. *Ann Microbiol*. 61:663–672.
- Bardhan, P.; Gohain, M.; Daimary, N.; Kishor, S.; Chattopadhyay, P.; Gupta, K. and Mandal, M. (2019). Microbial lipids from cellulolytic oleaginous fungus *Penicillium citrinum* PKB20 as a potential feedstock for biodiesel production. *Annals of Microbiology*. 69(11): 1135-1146
- Bellou, S.; Triantaphyllidou, I.E.; Aggeli, D.; Elazzazy, A.M.; Baeshen, M.N. and Aggelis, G. (2016). Microbial oils as food additives: Recent approaches for improving microbial oil production and its polyunsaturated fatty acid content. *Curr. Opin. Biotechnol*. 37: 24–35.
- Bezerra, M.A.; Santelli, R.E.; Oliveira, E.P.; Villar, L.S. and Escalera, L.A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*. 76 (5): 965–977.

- Carvalho, A.K.F.; Bento, H.B.; Izário Filho, H.J. and de Castro, H.F. (2018). Approaches to convert *Mucor circinelloides* lipid into biodiesel by enzymatic synthesis assisted by microwave irradiations. *Renew Energy*. 125:747–754.
- Chebbi, H.; Leiva-candia, D.; Carmona-cabello, M. and Jaouani A. (2019). Industrial Crops & Products Biodiesel production from microbial oil provided by oleaginous yeasts from olive oil mill wastewater growing on industrial glycerol. *Ind Crop Prod*. 139:111 - 535.
- Dong, T.; Knoshaug, E. P., Pienkos, P. T. and Laurens, L. M. (2016). Lipid recovery from wet oleaginous microbial biomass for biofuel production: a critical review. *Appl. Energy*. 177: 879–895.
- Economou, Ch.; Aggelis, N.; Pavlou, G. and Vayenas, S. (2011). Single cell oil production from rice hulls hydrolysate. *Bioresour Technol*. 102:9737–9742.
- Enshaeieh, M.; Nahvi, I. and Madani, M. (2014). Improving microbial oil production with standard and native oleaginous yeasts by using Taguchi design. *Int J Environ Sci Technol*. 11(3):597–604.
- Fawzy, M.A.; Darwish, H.; Alharthi, S.; Al-Zaban, M.I.; Noureldeen, A. and Hassan, S.H. (2022). Process optimization and modeling of Cd²⁺ biosorption onto the free and immobilized *Turbinaria ornata* using Box–Behnken experimental design. *Sci. Rep*. 12: 3256.
- Fazili, A.B.A.; Shah, A.M.; Zan, X.; Naz, T.; Nosheen, S.; Nazir, Y.; Ullah, S.; Zhang, H. and Song, Y. (2022). *Mucor circinelloides*: A model organism for oleaginous fungi and its potential applications in bioactive lipid production. *Microb. Cell Fact*. 21: 29.
- Folch, J.; Lees, M. and Sloane Stanley, G. H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem*. 226: 497–509.
- González-Fernández, M.J.; Ortea, I. and Guil-Guerrero, J.L. (2020). α -Linolenic and γ -linolenic acids exercise differential antitumor effects on HT-29 human colorectal cancer cells. *Toxicol. Res*.9: 474–483.
- Huang, C.; Luo, M.T.; Chen, X.F.; Qi, G.X.; Xiong, L.; Lin, X.Q. and Chenn, X.D. (2017). Combined “de novo” and “ex novo” lipid fermentation in a mix-medium of corncob acid hydrolysate and soybean oil by *Trichosporon dermatis*. *Biotechnology for Biofuels* 10: 1–11.

- Jiru, T.; Groenewald, M.; Pohl, C.; Stetn, L.; Kiggundu, N. and Abate D. (2017). Optimization of cultivation conditions for biotechnological production of lipid by *Rhodotorula kratochvilovae*(syn. *Rhodospiridium kratochvilovae*) SY89 for biodiesel preparation. *Biotech.* 7:1-1
- Kalampounias, G.; Gardeli, C.; Alexis, S.; Anagnostopoulou, E.; Androutsopoulou, T.; Dritsas, P.; Aggelis, G.; Papanikolaou, S. and Katsoris, P. (2024). Poly-unsaturated fatty acids (PUFAs) from *Cunninghamella elegans* grown on glycerol induce cell death and increase intracellular reactive oxygen species. *J. Fungi.* 10, 130. <https://doi.org/10.3390/jof10020130>
- Kumar, S. P. and Banerjee, R. (2013). Optimization of lipid enriched biomass production from oleaginous fungus using response surface methodology. *Indian Journal of Experimental Biology.* 51:979-983.
- Lei, Y.; Wang, X.; Sun, S.; He, B.; Sun, W.; Wang, K.; Chen, Z.; Guo, Z. and Li, Z. (2024). A review of lipid accumulation by oleaginous yeasts: Culture mode. *Science of the Total Environment.* 919: 170385.
- Leong, W.H.; Lim, J.W.; Lam, M.K.; Uemura, Y. and Ho, Y.C. (2018). Third generation biofuels: A nutritional perspective in enhancing microbial lipid production. *Renew Sustain Energy Rev.* 91:950–961.
- Lewis, T.; Nichols, P.D. and McMeekin, T.A. (2000). Evaluation of extraction methods for recovery of fatty acids from lipid-producing micro heterotrophs. *Journal of Microbiological Methods.*43: 107–116.
- Li, Q.; Du, W. and Liu, D. (2008). Perspectives of microbial oils for biodiesel production. *Appl Microbiol Biot Applied.* 80: 749.
- Lin, Y.; Wang, J.; Wang, X. and Sun, X. (2011). Optimization of butanol production from corn straw hydrolysate by *Clostridium acetobutylicum* using response surface method. *Chin. Sci. Bull.* 56 (14):1422–1428.
- Madani, M.; Enshaeieh, M. and Abdoli, A. (2017). Single cell oil and its application for biodiesel production. *Process Safety and Environmental Protection.* 111: 747–756.
- Manikan, V.; Kalil, M.S. and Hamid, A.A. (2015). Response surface optimization of culture medium for enhanced docosaheaxenoic acid production by a Malaysian thraustochytrid. *Sci. Rep.* 5: 8611. DOI:10.1038/srep08611.

- Mhlongo, S. I.; Ezeokoli, O. T.; Roopnarain, A.; Ndaba, B.; Sekoai, P. T.; Habimana, O. and Pohl, C. H. (2021). The Potential of Single-Cell Oils Derived From Filamentous Fungi as Alternative Feedstock Sources for Biodiesel Production. *Frontiers in Microbiology*. 12. 637381.
- Mironov, A. A.; Nemashkalov, V. A.; Stepanova, N. N.; Kamzolova, S. V.; Rymowicz, W. and Morgunov, I. G. (2018). The effect of pH and temperature on arachidonic acid production by glycerol-grown *Mortierella alpina* NRRLA-10995. *Fermentation*. 4:17.
- Moubasher, M.H.; Hamouda, M.S.; Tammam, AM. and Tahoun, I.F. (2016). Biodiesel production from agricultural and agro industrial wastes by *Fusarium oxysporum*. *Egypt. J. Bot.* 56(3): 733–751.
- Nasr, M.; Nahvi, I.; Keyhanfar, M. and Mirbagheri, M. (2017). The effect of carbon and nitrogen sources on the fatty acids profile of *Mortierella vinacea*. *Biological Journal of Microorganisms*. 5:1–8.
- Nisha, A. and Venkateswaran, G. (2011). Effect of culture variables on mycelial arachidonic acid production by *Mortierella alpina*. *Food and Bioprocess Technology*. 4:232–240.
- Nisha, A.; Muthukumar, S. P. and Venkateswaran, G. (2009). Safety evaluation of arachidonic acid rich *Mortierella alpine* biomass in albino rats-a sub chronic study. *Regulatory Toxicology and Pharmacology*. 53:186–194.
- Ochsenreither, K.; Glück, C.; Stressler, T.; Fischer, L. and Syldatk, C. (2016). Production strategies and applications of microbial single cell oils. *Front Microbiol.* 7: 1539.
- Papanikolaou, S. and Aggelis, G. (2019). Sources of microbial oils with emphasis to *Mortierella (Umbelopsis) isabellina* fungus. *World J. Microbiol. Biotechnol.* 35: 63.
- Papanikolaou, S.; Galiotou-Panayotou, M.; Fakas, S.; Komaitis, M. and Aggelis, G. (2007). Lipid production by oleaginous *Mucorales* cultivated on renewable carbon sources. *European Journal of Lipid Science and Technology*. 109(11): 1060-1070.
- Ramírez-Castrillón, M.; Jaramillo-García, V.P.; Rosa, P.D.; Landell, M.F.; Vu, D.; Fabricio, M.F.; Ayub, M.A.; Robert, V.; Henriques, J.A. and Valente, P. (2017). The oleaginous yeast *Meyerozyma guilliermondii*

- BI281A as a new potential biodiesel feedstock: selection and lipid production optimization. *Frontiers in microbiology*. 8: 1776.
- Ranjan, A.; Mayank, R. and Moholkar, V. (2013b). Process optimization for butanol production from developed rice straw hydrolysate using *Clostridium acetobutylicum* MTCC 481 strain. *Biomass Convers. Biorefin.* 3 (2):143–155.
- Ratledge, C. (1994). Yeasts, moulds, algae and bacteria as sources of lipids. In *Technological Advances in Improved and Alternative Sources of Lipids*; Kamel, B.S., Kakuda, Y., Eds.; Springer: Boston, MA, USA.: 235–291.
- Ratledge, C. and Wynn, J.P. (2002). The biochemistry and molecular biology of lipid accumulation in oleaginous microorganisms. In *Advances in Applied Microbiology*; Academic Press: Cambridge, MA, USA. 51: 1–52.
- Robles-Iglesias R.; Naveira-Pazos, C.; Fernández-Blanco, C.; Veiga, M.C. and Kennes, C. (2023). Factors affecting the optimisation and scale-up of lipid accumulation in oleaginous yeasts for sustainable biofuels production. *Renew. Sustain. Energy Rev.* 171: 113043.
- Rossi, M.; Amaretti, A.; Raimondi, S. and Leonardi, A. (2011). Getting lipids for biodiesel production from oleaginous fungi. *Feedstocks and Processing Technologies*. (InTech, Italy). 71.
- Ruan, Z.; Zanotti, M.; Archer, S.; Liao, W. and Liu, Y. (2014) . Oleaginous fungal lipid fermentation on combined acid- and alkali-pretreated corns over hydrolysate for advanced biofuel production. *Bioresource Technology*. 163: 12–17.
- Sheekh, M.M.; Gheda, S.F.; El-Sayed, A.E.B.; Abo Shady, A.M.; El-Sheikh, M.E. and Schagerl, M. (2019). Outdoor cultivation of the green microalga *Chlorella vulgaris* under culture stress conditions as a feedstock for biofuel. *Environ. Sci. Pollut. Res.* 26(18): 18520-18532.
- Shoab, A.; Bhran, A.; Rasmey, A. H. and Mikky, Y. (2018). Optimization of cultural conditions for lipid accumulation by *Aspergillus wentii* Ras101 and its transesterification to biodiesel: application of response surface methodology. *Biotech*, 8:1-11.
- Shrivastava, A.; Bajaj, I.; Saudagar, P. and Singhal, R. (2008). Media optimization for the production of γ -linolenic acid by *Cunninghamella*

- echinulata var. elegans MTCC 552 using response surface methodology. *Int J Food Eng.* 4:1-32.
- Singh, R.; Bishnoi, N.R. and Kirrolia, A. (2013). Evaluation of *Pseudomonas aeruginosa* an innovative bioremediation tool in multi metals ions from simulated system using multi response methodology. *Bioresour Technol.* 138: 222–234.
- Sukrutha, S.; Adamechova, K.; Savitha, J. and Certik M. (2014). Optimization of Physiological Growth Conditions for Maximal Gamma- linolenic Acid Production by *Cunninghamella blakesleeana* JSK2. *Journal of the American Oil Chemists' Society* .91:1507-1513.
- Xia, C.; Zhang, J.; Zhang, W. and Hu, B. (2011). A new cultivation method for microbial oil production: cell pelletization and lipid accumulation by *Mucor circinelloides*. *Biotechnol. Biofuels.* 4:15.
- Varma, P. S.; Chandrasekharan, S.; Venkateswaran, G.; Rajendran, S.; Mallapureddy, K. K.; Pandey, A. and Parameswaran, B. (2018). Optimization of Process Parameters for the Production of γ -Linolenic Acid by *Cunninghamella elegans* CFR C07 in Submerged Fermentation. *Food Technology and Biotechnology.* 56(1): 96.
- Venkata, S.G. and Venkata, M.S. (2014). Lipid accumulation for biodiesel production b oleaginous fungus *Aspergillus awamori*: influence of critical factors. *Fuel* 116: 509–515.
- Xing, D.; Wang, H.; Pan, A.; Wang, J. and Xue, D. (2012). Assimilation of corn fier hydrolysates and lipid accumulatonby *Mortierella isabellina*. *Biomass Bioenergy.* 39: 494-501.
- Youssef, G.; Elrefaey, A. and El-Assar, S. (2020). Potential Assessment of Oleaginous Fungi for Sustainable Biodiesel Production: Screening, Identification and Lipid Production Optimization. *Egyption journal of botany.* 1-14.
- Zheng, Z. M.; Hu, Q. L.; Hao, J.; Xu, F.; Guo, N. N.; Sun, Y. and Liu, D.H. (2008). Statistical optimization of culture conditions for 1,3-propanediol by *Klebsiella pneumoniae* AC 15 via central composite design. *Bioresour. Technol.* 99 (5), 1052–1056.